

Cefavex

CEFTIOFUR 50mg/ml



Suspension for injection for pigs and cattle



Injektionssuspension für Schweine und Rinder



Suspensión inyectable para porcino y bovino



Szuszpenziós injekció sertések és szarvasmarhák részére A.U.V.



Sospensione iniettabile per suini e bovini.



Zawiesina do wstrzykiwań dla świń i bydła



Suspensão injectável para suínos e bovinos



инжекционна суспензия за свине и говеда.



Suspensie injectabila pentru porcine și bovine



ΕΝΕΣΙΜΟ ΕΝΑΙΩΡΗΜΑ ΓΙΑ ΧΟΪΡΟΥΣ ΚΑΙ ΒΟΟΕΙΔΗ



s.p.[®] veterinaria

Cefavex 50 mg/ml

Suspension for injection for pigs and cattle

COMPOSITION

One ml of suspension contains:
Ceftiofur (as hydrochloride) 50,0 mg

TARGET SPECIES AND INDICATIONS FOR USE

Pigs:

For the treatment of bacterial respiratory disease associated with *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *Streptococcus suis*.

Cattle:

For the treatment of bacterial respiratory disease associated with *Mannheimia haemolytica* (previously *Pasteurella haemolytica*), *Pasteurella multocida* and *Histophilus somni* (previously *Haemophilus somnus*).

For the treatment of acute interdigital necrobacillosis (panaritium, foot rot), associated with *Fusobacterium necrophorum* and *Bacteroides melaninogenicus* (*Porphyromonas asaccharolytica*).

For treatment of the bacterial component of acute post-partum (puerperal) metritis within 10 days after calving associated with *Escherichia coli*, *Arcanobacterium pyogenes* and *Fusobacterium necrophorum*, sensitive to ceftiofur. The indication is restricted to cases where treatment with another antimicrobial has failed.

ADMINISTRATION ROUTE AND DOSE

Read the package leaflet before use.

Pigs:

3 mg ceftiofur /kg b.w./day for 3 days via intramuscular route, i.e. 1 ml/16 kg b.w. at each injection.

Cattle:

Respiratory disease: 1 mg ceftiofur /kg b.w./day for 3 to 5 days by subcutaneous injection, i.e. 1 ml/50 kg b.w. at each injection.
Acute interdigital necrobacillosis: 1 mg/kg b.w./day for 3 days by subcutaneous injection, i.e. 1 ml/50 kg b.w. at each injection.
Acute post-partum metritis within 10 days after calving: 1 mg/kg b.w./day for 5 consecutive days by subcutaneous injection, i.e. 1 ml/50 kg b.w. at each injection.

For detailed information on Pharmacodynamics and Pharmacokinetics, contact us through email or at our website.

WITHDRAWAL PERIOD

Pigs:

Meat and offal: 5 days.

Cattle:

Meat and offal: 8 days.

Milk: zero hours

PRESENTATION: 100 ml vials.

Notice: The product information provided is intended for informational purpose only. Please read the package insert of the product for proper use. Note that the label wording may be subdue to the local regulations of the country of destination.



s.p.® veterinaria, s.a.

Ctra. Reus-Vinyols Km. 4,1 • P.O. BOX 60 • 43330 RIUDOMS (Tarragona) SPAIN

Tel: +34 977 850 170 • Fax: +34 977 850 405

internacional@spveterinaria.com • www.spveterinaria.com