· ORAL SOLUTION ·

SOLUTION FOR USE IN DRINKING WATER FOR PIGS 300 mg/ml



No fever No ulcers No pain

Piresol® 300 mg/ml

SWINE: SYMPTOMATIC TREATMENT OF FEVER. 30 mg paracetamol / kg b.w.







Quick absorption. Effective plasma concentrations are obtained 10 min after ingestion, reaching maximum levels at 90 min.



Bioavailability is 81%, no plasma protein binding.

Quick elimination rate, no tissue accumulation.

STABILITY

Maintains a stable concentration in drinking water, superior to 95% of the initial concentration during 24 hours. The type of water quality does not affect this 24 h stability.



	% ULCERS	SEVERITY
Paracetamol (30 mg/kg b.w.)	0	-
ASA (50 ma/ka b.w.)	20	Superficial lesions

Does not alter the blood coagulation process

Large security margin

(5 times the dose and double administration time).

PARACETAMOL is a paraminophenol derivative with analgesic and antipyretic properties. Its antipyretic effect may be explained by its ability to inhibit brain cyclo-oxygenases. Paracetamol is a weak inhibitor of COX-1 synthesis, it has no gastro-intestinal side effects and has no effect on platelet-aggregation.

Composition:

Paracetamol300)	mg
Excipient c.s.p.	1	ml

Target species and indications:

Swine: Symptomatic treatment of fever.


Administration way and dosage:

Oral, through drinking water.

Swine: 30 mg de paracetamol/kg p.v./day, during 5 days. Equivalent, approx., to 1 L **PIRESOL 300 mg/ml** in 1000 L of drinking water.

Withdrawal time: 0 days. **Presentation:** 1 and 5 L. packs.

EFFICACY





Ctra. Reus-Vinyols Km. 4.1 • 43330 RIUDOMS (Tarragona) SPAIN Tel. +34 977 850 170* • Fax +34 977 850 405 • P.O. BOX 60 www.spveterinaria.com